

Program info:

Thursday, June 13, 2024
Start Time: 8:00 PM ET

Virtual Program

Once registered, you'll be emailed a link to access the virtual event



Speaker info:

David Fleischer, MD
Section Head of Allergy & Immunology
Children's Hospital Colorado Aurora, CO

Join us to learn about clinical efficacy and safety data for XOLAIR in IgE-mediated food allergy.

This program will include information on the OUtMATCH trial.

For more information, please contact:

- Michele Romesberg
- (814) 242-1940
- romesbem@gene.com

To register for this event, please visit:

- genentechsvp.com and enter CM45548
- Please RSVP by Wednesday, 6/12/2024

Minnesota, Vermont, and Federal Entities (eg, the Department of Defense and the Department of Veterans Affairs) have restrictions on receiving in-kind benefits (eg, meals, valet parking) at company-sponsored events. You are accountable for understanding such restrictions and complying with them. If you are licensed in or affiliated with any of these states or federal agencies, Genentech policies may restrict you from consuming any portion of the Genentech-sponsored meal at this program or from receiving any other in-kind benefit from Genentech (eg, valet parking) in connection with the program.

When you RSVP, please indicate whether you will accept or opt out of Genentech's in-kind benefits (eg, meals, valet parking) at the program. If you choose to opt out, you may either pay for the meal and parking on your own or not consume anything at the program.

For all program attendees who receive Genentech's in-kind benefits at this program, Genentech will report the attendee's name and the value received as required by federal and state disclosure laws (for more information on the federal law, please visit sunshine.gene.com).

The meal value reported may vary by event location and be up to \$150 per person (exceptions may apply).

INDICATION

XOLAIR® (omalizumab) is indicated for the reduction of allergic reactions (Type I), including anaphylaxis, that may occur with accidental exposure to one or more foods in adult and pediatric patients aged 1 year and older with IgE-mediated food allergy.

XOLAIR is to be used in conjunction with food allergen avoidance.

Limitations of Use: XOLAIR is not indicated for the emergency treatment of allergic reactions, including anaphylaxis.

IMPORTANT SAFETY INFORMATION

WARNING: Anaphylaxis

Anaphylaxis presenting as bronchospasm, hypotension, syncope, urticaria, and/or angioedema of the throat or tongue, has been reported to occur after administration of XOLAIR. Anaphylaxis has occurred as early as after the first dose of XOLAIR, but also has occurred beyond 1 year after beginning regularly administered treatment. Because of the risk of anaphylaxis, initiate XOLAIR therapy in a healthcare setting and closely observe patients for an appropriate period of time after XOLAIR administration. Health care providers administering XOLAIR should be prepared to manage anaphylaxis which can be life-threatening. Inform patients of the signs and symptoms of anaphylaxis and instruct them to seek immediate medical care should symptoms occur. Selection of patients for self-administration of XOLAIR should be based on criteria to mitigate risk from anaphylaxis.

CONTRAINDICATIONS

XOLAIR is contraindicated in patients with a severe hypersensitivity reaction to XOLAIR or to any ingredient of XOLAIR.

WARNINGS AND PRECAUTIONS

Anaphylaxis: Anaphylaxis has been reported to occur after administration of XOLAIR in premarketing clinical trials and in postmarketing spontaneous reports. In premarketing clinical trials in patients for a different indication, anaphylaxis was reported in 3 of 3507 (0.1%) patients. Anaphylaxis occurred with the first dose of XOLAIR in two patients and with the fourth dose in one patient. The time to onset of anaphylaxis was 90 minutes after administration in two patients and 2 hours after administration in one patient.

A case-control study in asthma patients showed that, among XOLAIR users, patients with a history of anaphylaxis to foods, medications, or other causes were at increased risk of anaphylaxis associated with XOLAIR, compared to those with no prior history of anaphylaxis. In postmarketing spontaneous reports, the frequency of anaphylaxis attributed to XOLAIR use was estimated to be at least 0.2% of patients based on an estimated exposure of about 57,300 patients from June 2003 through December 2006. Approximately 60% to 70% of anaphylaxis cases have been reported to occur within the first three doses of XOLAIR, with additional cases occurring sporadically beyond the third dose.

IMPORTANT SAFETY INFORMATION (cont'd)

WARNINGS AND PRECAUTIONS (cont'd)

Initiate XOLAIR only in a healthcare setting equipped to manage anaphylaxis which can be life-threatening. Observe patients closely for an appropriate period of time after administration of XOLAIR, taking into account the time to onset of anaphylaxis seen in premarketing clinical trials and postmarketing spontaneous reports. Inform patients of the signs and symptoms of anaphylaxis, and instruct them to seek immediate medical care should signs or symptoms occur.

Once XOLAIR therapy has been established, administration of XOLAIR prefilled syringe or autoinjector outside of a healthcare setting by a patient or a caregiver may be appropriate for selected patients. Patient selection, determined by the healthcare provider in consultation with the patient, should take into account the pattern of anaphylaxis events seen in premarketing clinical trials and postmarketing spontaneous reports, as well as individual patient risk factors (e.g. prior history of anaphylaxis), ability to recognize signs and symptoms of anaphylaxis, and ability to perform subcutaneous injections with XOLAIR prefilled syringe or autoinjector with proper technique according to the prescribed dosing regimen and Instructions for Use.

Discontinue XOLAIR in patients who experience a severe hypersensitivity reaction.

Malignancy: Malignant neoplasms were observed in 20 of 4127 (0.5%) XOLAIR-treated patients compared with 5 of 2236 (0.2%) control patients in clinical studies of adults and adolescents (≥ 12 years of age) for a different indication and other allergic disorders. The observed malignancies in XOLAIR-treated patients were a variety of types, with breast, non-melanoma skin, prostate, melanoma, and parotid occurring more than once, and five other types occurring once each. The majority of patients were observed for less than 1 year. The impact of longer exposure to XOLAIR or use in patients at higher risk for malignancy (e.g., elderly, current smokers) is not known.

A subsequent 5-year observational study of 5007 XOLAIR-treated and 2829 non-XOLAIR-treated adolescent and adult patients for a different indication found that the incidence rates of primary malignancies (per 1000 patient years) were similar in both groups (12.3 vs 13.0, respectively). Study limitations which include the observational study design, the bias introduced by allowing enrollment of patients previously exposed to XOLAIR (88%), enrollment of patients (56%) while a history of cancer or a premalignant condition were study exclusion criteria, and the high study discontinuation rate (44%) preclude definitively ruling out a malignancy risk with XOLAIR.

Fever, Arthralgia, and Rash: In post-approval use, some patients have experienced a constellation of signs and symptoms, including arthritis/arthralgia, rash, fever, and lymphadenopathy with an onset 1 to 5 days after the first or subsequent injections of XOLAIR. These signs and symptoms have recurred after additional doses in some patients. Physicians should stop XOLAIR if a patient develops this constellation of signs and symptoms.

Parasitic (Helminth) Infection: Monitor patients at high risk of geohelminth infection while on XOLAIR therapy. Insufficient data are available to determine the length of monitoring required for geohelminth infections after stopping XOLAIR treatment.

Laboratory Tests: Due to formation of XOLAIR:IgE complexes, serum total IgE levels increase following administration of XOLAIR and may remain elevated for up to 1 year following discontinuation of XOLAIR. Do not use serum total IgE levels obtained less than 1 year following discontinuation to reassess the dosing regimen for IgE-mediated food allergy patients, because these levels may not reflect steady state free IgE levels.

Potential Medication Error Related to Emergency Treatment of Anaphylaxis

XOLAIR should not be used for the emergency treatment of allergic reactions, including anaphylaxis. In studies to simulate use, some patients and caregivers did not understand that XOLAIR is not intended for the emergency treatment of allergic reactions, including anaphylaxis. The safety and effectiveness of XOLAIR for emergency treatment of allergic reactions, including anaphylaxis, have not been established. Instruct patients that XOLAIR is for maintenance use to reduce allergic reactions, including anaphylaxis, while avoiding food allergens.

ADVERSE REACTIONS

IgE-Mediated Food Allergy: The most common adverse reactions ($\geq 3\%$ incidence in XOLAIR-treated patients) included: injection site reaction (15.5%) and pyrexia (6.4%). Safety data obtained from adults ($n=3$) in this trial was limited.

Injection Site Reactions: Injection site reactions occurred at a rate of 15.5% in XOLAIR-treated patients compared with 10.9% in placebo-treated patients. The types of injection site reactions included: urticaria, discomfort, erythema, pain, and rash. All injection site reactions were mild to moderate severity and none resulted in study discontinuation.

Injection Site Reactions in Healthy Adults: In an open label trial in healthy adults, in which the 300 mg/2 mL autoinjector was compared to the 300 mg/2 mL prefilled syringe, injection site reactions (e.g., induration, pain, erythema, hemorrhage, swelling, discomfort, bruising, hypoesthesia, edema, pruritus) were observed in 24% (16/66) of subjects treated with the autoinjector compared with 14% (9/64) of subjects treated with the prefilled syringe.

Cardiovascular and Cerebrovascular Events from Clinical Studies for a Different

Indication: A 5-year observational study was conducted in 5007 XOLAIR-treated and 2829 non-XOLAIR-treated patients ≥ 12 years of age for a different indication to evaluate the long term safety of XOLAIR, including the risk of malignancy. Similar percentages of patients in both cohorts were current (5%) or former smokers (29%). Patients had a mean age of 45 years and were followed for a mean of 3.7 years. More XOLAIR-treated patients were diagnosed with a severe form of the condition studied (50%) compared to the non-XOLAIR-treated patients (23%). A higher incidence rate (per 1000 patient-years) of overall cardiovascular and cerebrovascular serious adverse events (SAEs) was observed in XOLAIR-treated patients (13.4) compared to non-XOLAIR-treated patients (8.1). Increases in rates were observed for transient ischemic attack (0.7 vs 0.1), myocardial infarction (2.1 vs 0.8), pulmonary hypertension (0.5 vs 0), pulmonary embolism/venous thrombosis (3.2 vs 1.5), and unstable angina (2.2 vs 1.4), while the rates observed for ischemic stroke and cardiovascular death were similar among both study cohorts. The results suggest a potential increased risk of serious cardiovascular and cerebrovascular events in patients treated with XOLAIR, however the observational study design, the inclusion of patients previously exposed to XOLAIR (88% for a mean of 8 months), baseline imbalances in cardiovascular risk factors between the treatment groups, an inability to adjust for unmeasured risk factors, and the high study discontinuation rate (44%) limit the ability to quantify the magnitude of the risk.

Pregnancy: Data with XOLAIR use in pregnant women are insufficient to inform on drug associated risk.

You may report side effects to the FDA at (800) FDA-1088 or www.fda.gov/medwatch. You may also report side effects to Genentech at (888) 835-2555 or Novartis Pharmaceuticals Corporation at (888) 669-6682.

Please see accompanying full Prescribing Information, including Boxed WARNING and Medication Guide, for additional Important Safety Information.